

Publications

Dr Simon J. GEORGE

	[1] Spectroscopic characterization of ferredoxin I and ferredoxin II from <i>Desulfovibrio africanus</i> Hatchikian, E. C.; Cammack, R.; Patil, D. S.; Robinson, A. E.; Richards, A. J. M.; George, S. J. ; Thomson, A. J. <i>Biochim. Biophys. Acta</i> 1984 , 784, 40-47.
1984	[2] <i>Azotobacter chroococcum</i> 7Fe ferredoxin - 2 pH dependent forms of the reduced 3Fe cluster and its conversion to a 4Fe cluster George, S. J. ; Richards, A. J. M.; Thomson, A. J.; Yates, M. G. <i>Biochem. J.</i> 1984 , 224, 247-251.
1985	[3] A study of one of the iron-sulfur clusters in oxidized hydrogenase from <i>Megasphaera elsdenii</i> by Magnetic Circular Dichroism Spectroscopy Thomson, A. J.; George, S. J. ; Richards, A. J. M.; Robinson, A. E.; Grande, H. J.; Veeger, C.; van Dijk, C. <i>Biochem. J.</i> 1985 , 227, 333-336.
1986	[4] Magnetic circular dichroism of iron-sulphur proteins George, S. J. , PhD Thesis, University of East Anglia, 1986 .
1988	[5] Direct electrochemistry in the characterization of redox proteins - novel properties of <i>Azotobacter</i> 7Fe ferredoxin Armstrong, F. A.; George, S. J. ; Thomson, A. J.; Yates, M. G. <i>Febs Lett.</i> 1988 , 234, 107-110. [6] X-ray crystallography and the spectroscopic imperative - the story of the [3Fe-4S] clusters George, G. N. ; George, S. J. <i>Trends in Biochemical Sciences</i> 1988 , 13, 369-370. [7] Redox properties of <i>Azotobacter</i> 7Fe ferredoxin Armstrong, F. A.; George, S. J. ; Thomson, A. J.; Yates, M. G. <i>Biochem. Soc. Trans.</i> 1988 , 16, 840-842. [8] NMR, EPR and magnetic-CD studies of cytochrome-f - identity of the heme axial ligands Rigby, S. E. J.; Moore, G. R.; Gray, J. C.; Gadsby, P. M. A.; George, S. J. ; Thomson, A. J. <i>Biochem. J.</i> 1988 , 256, 571-577.
1989	[9] Electrochemical and spectroscopic characterization of the 7Fe form of ferredoxin III from <i>Desulfovibrio africanus</i> Armstrong, F. A.; George, S. J. ; Cammack, R.; Hatchikian, E. C.; Thomson, A. J. <i>Biochem. J.</i> 1989 , 264, 265-273. [10] Electrochemical and spectroscopic characterization of the conversion of the 7Fe into the 8Fe Form of ferredoxin III from <i>Desulfovibrio africanus</i> - identification of a [4Fe-4S] cluster with one non-cysteine ligand George, S. J. ; Armstrong, F. A.; Hatchikian, E. C.; Thomson, A. J. <i>Biochem. J.</i> 1989 , 264, 275-284. [11] Evidence for reversible multiple redox transformations of [3Fe-4S] clusters Armstrong, F. A.; Butt, J. N.; George, S. J. ; Hatchikian, E. C.; Thomson, A. J. <i>Febs Lett.</i> 1989 , 259, 15-18.
1990	[12] Electrochemical and spectroscopic studies of ferredoxins Thomson, A. J.; George, S. J. ; Armstrong, F. A.; Hatchikian, E. C.; Yates, M. G. <i>Pure and Applied Chemistry</i> 1990 , 62, 1071-1074. [13] Cytochrome-c" isolated from <i>Methylophilus methylotrophus</i> - an example of bis-Histidine coordinated Fe ³⁺ heme, with near-perpendicular orientation of the ligands Berry, M. J.; George, S. J. ; Thomson, A. J.; Santos, H.; Turner, D. L. <i>Biochem. J.</i> 1990 , 270, 413-417.
1991	[14] A comparative study of the low-temperature Magnetic Circular Dichroism spectra of ferredoxins containing [4Fe-4S] and [4Fe-4Se] clusters George, S. J. ; Thomson, A. J.; Crabtree, D. E.; Meyer, J.; Moulis, J. M. <i>New J. Chem.</i> 1991 , 15, 455-465. [15] Iron-sulfur clusters in electron-transfer, catalysis and control Thomson, A. J.; Breton, J.; George, S. J. ; Butt, J. N.; Armstrong, F. A.; Hatchikian, E. C. <i>Biochem. Soc. Trans.</i> 1991 , 19, 594-599. [16] Fluorescence XAFS detectors for 0.3 - 20 keV. Cramer, S. P.; George, S. J. ; Kraner, H.; Rogers, L.; Rescia, S.; Radeka, V.; Yocom, M.; Colaresi, J.; Tench, O.; Mullins, O. C., in <i>X-ray Absorption Fine Structure</i> ; Hasnain, S. S., Ed.; Ellis Harwood: New York, 1991 , pp. 640-645.

	[17] Investigation of metal-ion uptake reactivities of [3Fe-4S] clusters in proteins - Voltammetry of coadsorbed ferredoxin-aminocyclitol films at graphite electrodes and spectroscopic identification of transformed clusters Butt, J. N.; Armstrong, F. A.; Breton, J.; George, S. J. ; Thomson, A. J.; Hatchikian, E. C. <i>J. Am. Chem. Soc.</i> 1991 , <i>113</i> , 6663-6670.
	[18] Identification and characterization of zinc binding sites in protein kinase C Hubbard, S. R.; Bishop, W. R.; Kirschmeier, P.; George, S. J. ; Cramer, S. P.; Hendrickson, W. A. <i>Science</i> 1991 , <i>254</i> , 1776-1779.
1992	[19] L-edge x-ray absorption spectroscopy of <i>Pyrococcus furiosus</i> rubredoxin George, S. J. ; van Elp, J.; Chen, J.; Ma, Y.; Chen, C. T.; Park, J. B.; Adams, M. W. W.; Searle, B. G.; Degroot, F. M. F.; Fuggle, J. C.; Cramer, S. P. <i>J. Am. Chem. Soc.</i> 1992 , <i>114</i> , 4426-4427.
	[20] Soft x-ray spectroscopy of metalloproteins using fluorescence detection Cramer, S. P.; Chen, J.; George, S. J. ; van Elp, J.; Moore, J.; Tench, O.; Colaresi, J.; Yocom, M.; Mullins, O. C.; Chen, C. T. <i>Nucl. Instrum. Meth. A</i> 1992 , <i>319</i> , 285-289.
	[21] Nitrogen chemical structure in DNA and related molecules by x-ray absorption spectroscopy Mitra-Kirtley, S.; Mullins, O. C.; Chen, J.; van Elp, J.; George, S. J. ; Chen, C. T.; Ohalloran, T.; Cramer, S. P. <i>Biochim. Biophys. Acta</i> 1992 , <i>1132</i> , 249-254.
	[22] Determination of the nitrogen chemical structures in petroleum asphaltenes using XANES spectroscopy Mitra-Kirtley, S.; Mullins, O. C.; van Elp, J.; George, S. J. ; Chen, J.; Cramer, S. P. <i>J. Am. Chem. Soc.</i> 1993 , <i>115</i> , 252-258.
	[23] Variable temperature magnetic circular dichroism Thomson, A. J.; Cheesman, M. R.; George, S. J. , in <i>Meth. Enzymol.</i> , 1993 , <i>225</i> , 199-232.
	[24] Copper L-edge spectral studies - a direct experimental probe of the ground-state covalency in the blue copper site in plastocyanin George, S. J. ; Lowery, M. D.; Solomon, E. I.; Cramer, S. P. <i>J. Am. Chem. Soc.</i> 1993 , <i>115</i> , 2968-2969.
	[25] Iron EXAFS of <i>Azotobacter vinelandii</i> nitrogenase Mo-Fe and V-Fe proteins Chen, J.; Christiansen, J.; Tittsworth, R. C.; Hales, B. J.; George, S. J. ; Coucouvanis, D.; Cramer, S. P. <i>J. Am. Chem. Soc.</i> 1993 , <i>115</i> , 5509-5515.
1993	[26] Extended x-ray absorption fine structure and L-edge spectroscopy of nitrogenase molybdenum-iron protein Chen, J.; Christiansen, J.; George, S. J. ; Tittsworth, R. C.; Hales, B. J.; Alahmad, S.; Coucouvanis, D.; Campobasso, N.; Bolin, J.; Cramer, S. P., in <i>Molybdenum Enzymes, Cofactors, and Model Systems</i> ; Steifel, E. I., Coucouvanis, D. and Newton, W. E., Ed.; American Chemical Society: Washington D.C., 1993 , pp. 231-242.
	[27] X-ray absorption spectroscopy of nitrogenase MoFe protein Chen, J.; Christiansen, J.; George, S. J. ; Tittsworth, R. C.; Hales, B. J.; Campobasso, N.; Bolin, J.; Cramer, S. P., in <i>Applications of Synchrotron Radiation Techniques to Materials Science</i> ; Perry, S. L., D., S. N., Stockbauer, R. L., d'Amico, K. L. and Terminello, L. J., Ed.; Materials Research Society; Vol. 307, 1993 .
	[28] Soft x-ray magnetic circular dichroism on a paramagnetic bioinorganic system Vanelp, J.; George, S. J. ; Peng, G.; Searle, B. G.; Zhou, Z. H.; Adams, M. W. W.; Chen, C. T.; Cramer, S. P. <i>Proceedings SPIE</i> 1993 , <i>2010</i> , 181
	[29] Soft x-ray magnetic circular dichroism - a probe for studying paramagnetic bioinorganic systems Vanelp, J.; George, S. J. ; Chen, J.; Peng, G.; Chen, C. T.; Tjeng, L. H.; Meigs, G.; Lin, H. J.; Zhou, Z. H.; Adams, M. W. W.; Searle, B. G.; Cramer, S. P. <i>Proc. Natl. Acad. Sci. U.S.A.</i> 1993 , <i>90</i> , 9664-9667.
1994	[30] Soft x-ray absorption and x-ray magnetic circular dichroism in biology George, S. J. ; van Elp, J.; Chen, J.; Peng, G.; Mitra-Kirtley, S.; Mullins, O. C.; Cramer, S. P., in <i>Synchrotron Radiation in Biosciences</i> ; Chance, B., Ed.; Clarendon Press: Oxford, 1994 , pp. 313-332.
1995	[31] Biological x-ray spectroscopy on 3rd-generation synchrotron radiation sources Ralston, C. Y.; Chen, J.; Peng, G.; George, S. J. ; van Elp, J.; Cramer, S. P. <i>Physica B</i> 1995 , <i>209</i> , 203-208.

	[32] Identification of the iron-sulfur clusters in a ferredoxin from the archaeon <i>Sulfolobus acidocaldarius</i> - evidence for a reduced [3Fe-4S] cluster with pH dependent electronic properties Breton, J. L.; Duff, J. L. C.; Butt, J. N.; Armstrong, F. A.; George, S. J. ; Petillot, Y.; Forest, E.; Schafer, G.; Thomson, A. J. <i>Eur. J. Biochem.</i> 1995 , 233, 937-946.
1996	[33] Manganese L-edge X-ray absorption spectroscopy of manganese catalase from <i>Lactobacillus plantarum</i> and mixed valence manganese complexes Grush, M. M.; Chen, J.; Stemmler, T. L.; George, S. J. ; Ralston, C. Y.; Stibrany, R. T.; Gelasco, A.; Christou, G.; Gorun, S. M.; Penner-Hahn, J. E.; Cramer, S. P. <i>J. Am. Chem. Soc.</i> 1996 , 118, 65-69.
	[34] Fluorescence Detected Soft XAS and MCD - Applications to Bioinorganic Systems Cramer, S. P.; Peng, G.; Christiansen, J.; Chen, J.; van Elp, J.; George, S. J. ; Young, A. T. <i>J. Electron Spectrosc.</i> 1996 , 78, 225-229.
	[35] Anionic tobacco peroxidase is active at extremely low pH: Veratryl alcohol oxidation with a pH optimum of 1.8 Gazarian, I. G.; Lagrimini, L. M.; George, S. J. ; Thorneley, R. N. F. <i>Biochem. J.</i> 1996 , 320, 369-372.
	[36] Iron L-edge X-ray absorption spectroscopy of myoglobin complexes and photolysis products Wang, H. X.; Peng, G.; Miller, L. M.; Scheuring, E. M.; George, S. J. ; Chance, M. R.; Cramer, S. P. <i>J. Am. Chem. Soc.</i> 1997 , 119, 4921-4928.
1997	[37] Time-resolved binding of carbon monoxide to nitrogenase monitored by stopped-flow infrared spectroscopy George, S. J. ; Ashby, G. A.; Wharton, C. W.; Thorneley, R. N. F. <i>J. Am. Chem. Soc.</i> 1997 , 119, 6450-6451.
	[38] Salicylic acid is a reducing substrate and not an effective inhibitor of ascorbate peroxidase Kvaratskhelia, M.; George, S. J. ; Thorneley, R. N. F. <i>J. Biol. Chem.</i> 1997 , 272, 20998-21001.
	[39] The structure of carbonyl horseradish peroxidase: spectroscopic and kinetic characterization of the carbon monoxide complexes of His-42 → Leu and Arg-38 → Leu mutants Rodriguez-Lopez, J. N.; George, S. J. ; Thorneley, R. N. F. <i>J. Biol. Inorg. Chem.</i> 1998 , 3, 44-52.
1999	[40] A novel, general method of analyzing magnetic circular dichroism spectra and magnetization curves of high-spin metal ions: Application to the protein oxidized rubredoxin, <i>Desulfovibrio gigas</i> Oganesyan, V. S.; George, S. J. ; Cheesman, M. R.; Thomson, A. J. <i>Journal of Chemical Physics</i> 1999 , 110, 762-777.
	[41] Reversible alkaline inactivation of lignin peroxidase involves the release of both the distal and proximal site calcium ions and bis-histidine co-ordination of the haem George, S. J. ; Kvaratskhelia, M.; Dilworth, M. J.; Thorneley, R. N. F. <i>Biochem. J.</i> 1999 , 344, 237-244.
	[42] Quantitation of metal ion and DNA junction binding to the Holliday junction endonuclease Cce1 Kvaratskhelia, M.; George, S. J. ; Cooper, A.; White, M. F. <i>Biochemistry</i> 1999 , 38, 16613-16619.
2000	[43] Time resolved infrared spectroscopy of functioning nitrogenase. Thorneley, R. N. F.; George, S. J. , in <i>Prokaryotic nitrogen fixation: a model system for analysis of a biological process.</i> ; E.W., T., Ed.; Horizon Scientific Press: Wymondham, UK., 2000 , pp. 81-99
	[44] Time-resolved infrared spectroscopy reveals a stable ferric heme-NO intermediate in the reaction of Paracoccus pantotrophus cytochrome cd(1) nitrite reductase with nitrite George, S. J. ; Allen, J. W. A.; Ferguson, S. J.; Thorneley, R. N. F. <i>J. Biol. Chem.</i> 2000 , 275, 33231-33237.
	[45] Stopped-Flow Infra-Red Spectroscopy of Carbon Monoxide Binding to Functioning Nitrogenase Thorneley, R. N. F.; Ashby, G. A.; George, S. J. , in <i>Nitrogen Fixation: From Molecules to Crop Productivity</i> ; Pedrosa, F. O., Hungria, M., Yates, G. and Newton, W. E., Ed.; Kluwer: The Netherlands, 2000 , pp. 39-40.
2001	[46] Stopped-flow infrared spectroscopy reveals a six-coordinate intermediate in the formation of the proximally bound five-coordinate NO adduct of cytochrome c' George, S. J. ; Andrew, C. R.; Lawson, D. M.; Thorneley, R. N. F.; Eady, R. R. , <i>J. Am. Chem. Soc.</i> 2001 , 123, 9683-9684.

	[47] Six- to five-coordinate heme-nitrosyl conversion in cytochrome c' and its relevance to guanylate cyclase Andrew, C. R.; George, S. J. ; Lawson, D. M.; Eady, R. R. <i>Biochemistry</i> 2002 , <i>41</i> , 2353-2360.
2002	[48] Transient FTIR spectroelectrochemical and stopped-flow detection of a mixed valence {Fe(I)-Fe(II)} bridging carbonyl intermediate with structural elements and spectroscopic characteristics of the di-iron sub-site of all-iron hydrogenase Razavet, M.; Borg, S. J.; George, S. J. ; Best, S. P.; Fairhurst, S. A.; Pickett, C. J., <i>Chem. Comm.</i> 2002 , 700-701.
	[49] The di-iron subsite of all-iron hydrogenase: Mechanism of cyanation of a synthetic {2Fe3S} - Carbonyl assembly George, S. J. ; Cui, Z.; Razavet, M.; Pickett, C. J., <i>Chem. Eur. J.</i> 2002 , <i>8</i> , 4037-4046.
	[50] A two-faced molecule offers NO explanation: the proximal binding of nitric oxide to haem Lawson, D. M.; Stevenson, C. E. M.; Andrew, C. R.; George, S. J. ; Eady, R. R. <i>Biochem. Soc. Trans.</i> 2003 , <i>31</i> , 553-557.
2003	[51] Stopped-flow Fourier transform infrared spectroscopy of nitromethane oxidation by the diiron(IV) intermediate of methane monooxygenase Muthusamy, M.; Ambundo, E. A.; George, S. J. ; Lippard, S. J.; Thorneley, R. N. F. <i>J. Am. Chem. Soc.</i> 2003 , <i>125</i> , 11150-11151.
	[52] Hydrogen-induced activation of the [NiFe]-hydrogenase from <i>Allochromatium vinosum</i> as studied by stopped-flow infrared spectroscopy Kurkin, S.; George, S. J. ; Thorneley, R. N. F.; Albracht, S. P. <i>J. Biochemistry</i> 2004 , <i>43</i> , 6820-6831.
2004	[53] Reactions of H-2, CO, and O-2 with active [NiFe]-Hydrogenase from <i>Allochromatium vinosum</i> . A stopped-flow infrared study George, S. J. ; Kurkin, S.; Thorneley, R. N. F.; Albracht, S. P. <i>J. Biochemistry</i> 2004 , <i>43</i> , 6808-6819.
	[54] X-ray magnetic circular dichroism - a high energy probe of magnetic properties Funk, T.; Deb, A.; George, S. J. ; Wang, H. X.; Cramer, S. P. <i>Coord. Chem. Rev.</i> 2005 , <i>249</i> , 3-30.
2005	[55] High-resolution X-ray emission spectroscopy of molybdenum compounds Doonan, C. J.; Zhang, L. M.; Young, C. G.; George, S. J. ; Deb, A.; Bergmann, U.; George, G. N.; Cramer, S. P. <i>Inorg. Chem.</i> 2005 , <i>44</i> , 2579-2581.
	[56] Normal mode analysis of FeCl4- and Fe2S2Cl42- via vibrational Mössbauer, resonance Raman, and FT-IR spectroscopies Smith, M. C.; Xiao, Y. M.; Wang, H. X.; George, S. J. ; Coucouvanis, D.; Koutmos, M.; Sturhahn, W.; Alp, E. E.; Zhao, J. Y.; Cramer, S. P. <i>Inorg. Chem.</i> 2005 , <i>44</i> , 5562-5570.
	[57] EPR and infrared spectroscopic evidence that a kinetically competent paramagnetic intermediate is formed when acetyl-coenzyme A synthase reacts with CO George, S. J. ; Seravalli, J.; Ragsdale, S. W. <i>J. Am. Chem. Soc.</i> 2005 , <i>127</i> , 13500-13501.
	[58] Reductive activation of nitrate reductases Field, S. J.; Thornton, N. P.; Anderson, L. J.; Gates, A. J.; Reilly, A.; Jepson, B. J. N.; Richardson, D. J.; George, S. J. ; Cheesman, M. R.; Butt, J. N. <i>Dalton Trans.</i> 2005 , 3580-3586.
	[59] Normal mode analysis of <i>Pyrococcus furiosus</i> rubredoxin via nuclear resonance vibrational spectroscopy (NRVS) and resonance Raman spectroscopy Xiao, Y. M.; Wang, H.; George, S. J. ; Smith, M. C.; Adams, M. W. W.; Jenney, F. E.; Sturhahn, W.; Alp, E. E.; Zhao, J. O.; Yoda, Y.; Dey, A.; Solomon, E. I.; Cramer, S. P. <i>J. Am. Chem. Soc.</i> 2005 , <i>127</i> , 14596-14606.
	[60] The advantages of soft X-rays and cryogenic spectrometers for measuring chemical speciation by X-ray spectroscopy Drury, O. B.; Friedrich, S.; George, S. J. ; Cramer, S. P. <i>Nucl. Instrum. Meth. A</i> 2006 , <i>559</i> , 728-730.
	[61] How nitrogenase shakes - Initial information about P-cluster and FeMo-cofactor normal modes from nuclear resonance vibrational Spectroscopy (NRVS) Xiao, Y. M.; Fisher, K.; Smith, M. C.; Newton, W. E.; Case, D. A.; George, S. J. ; Wang, H. X.; Sturhahn, W.; Alp, E. E.; Zhao, J. Y.; Yoda, Y.; Cramer, S. P. <i>J. Am. Chem. Soc.</i> 2006 , <i>128</i> , 7608-7612.

	[62] Spectroscopic and computational studies of reduction of the metal versus the tetrapyrrole ring of coenzyme F-430 from methyl-coenzyme M reductase Dey, M.; Kunz, R. C.; van Heuvelen, K. M.; Craft, J. L.; Horng, Y. C.; Tang, Q.; Bocian, D. F.; George, S. J. ; Brunold, T. C.; Ragsdale, S. W. <i>Biochemistry</i> 2006 , <i>45</i> , 11915-11933.
2007	[63] Identification of a Mo-Fe-S cluster on NifEN by Mo K-edge extended X-ray absorption fine structure George, S. J. ; Igarashi, R. Y.; Piamonteze, C.; Soboh, B.; Cramer, S. P.; Rubio, L. M. <i>J. Am. Chem. Soc.</i> 2007 , <i>129</i> , 3060-3061.
	[64] Activation of the cytochrome cd(1) nitrite reductase from <i>Paracoccus pantotrophus</i> - Reaction of oxidized enzyme with substrate drives a ligand switch at heme c van Wonderen, J. H.; Knight, C.; Oganesyan, V. S.; George, S. J. ; Zumft, W. G.; Cheesman, M. R. <i>J. Biol. Chem.</i> 2007 , <i>282</i> , 28207-28215.
	[65] The superconducting high-resolution soft X-ray spectrometer at the advanced biological and environmental X-ray facility Friedrich, S.; Drury, O. B.; George, S. J. ; Cramer, S. P. <i>Nucl. Instrum. Meth. A</i> 2007 , <i>582</i> , 187-189.
	[66] Protein voltammetry and spectroscopy: integrating approaches Male, L.; Marriott, S. J.; Berks, B. C.; Cheesman, M. R.; van Wonderen, J. H.; George, S. J. ; Butt, J. N. <i>Theor. Chem. Acc.</i> 2008 , <i>119</i> , 107-111.
2008	[67] X-ray photochemistry in iron complexes from Fe(0) to Fe(IV) - Can a bug become a feature? George, S. J. ; Fu, J.; Guo, Y.; Drury, O. B.; Friedrich, S.; Rauchfuss, T.; Volkers, P. I.; Peters, J. C.; Scott, V.; Brown, S. D.; Thomas, C. M.; Cramer, S. P. <i>Inorg. Chim. Acta</i> 2008 , <i>361</i> , 1157-1165.
	[68] EXAFS and NRVS reveal that NifB-co, a FeMo-co precursor, comprises a 6Fe core with an interstitial light atom George, S. J. ; Igarashi, R. Y.; Xiao, Y.; Hernandez, J. A.; Demuez, M.; Zhao, D.; Yoda, Y.; Ludden, P. W.; Rubio, L. M.; Cramer, S. P. <i>J. Am. Chem. Soc.</i> 2008 , <i>130</i> , 5673-5680.
	[69] Star-shaped molecule of Mn ^{II} ₄ O ₆ Core with a St = 10 high-spin State. A theoretical and experimental study with XPS, XMCD, and other magnetic methods. Khanra, S.; Kuepper, K.; Weyhermüller, T.; Prinz, M.; Raekers, M.; Voget, S.; Postnikov, A. V.; de Groot, F. M. F.; George, S. J. ; Coldea, M.; Neumann, M.; Chaudhuri, P. <i>Inorg. Chem.</i> 2008 , <i>47</i> , 4605-4617.
	[70] Molybdenum X-ray absorption edges from 200 - 20,000 eV, The benefits of soft x-ray spectroscopy for chemical speciation George, S. J. ; Drury, O. B.; Fu, J.; Friedrich, S.; Doonan, C. J.; George, G. N.; White, J. M.; Young, C. G.; Cramer, S. P. <i>J. Inorg. Biochem.</i> , 2009 , <i>103</i> , 157-167
2009	[71] Molybdenum Trafficking for Nitrogen Fixation Hernandez, J. A.; George, S. J. ; Rubio, L. M. <i>Biochemistry</i> , 2009 , <i>48</i> , 9711-9721.
	[72] A Star-Shaped Heteronuclear (CrMn3II)-Mn-III Species and Its Precise Electronic and Magnetic Structure: Spin Frustration Studied by X-Ray Spectroscopic, Magnetic, and Theoretical Methods Prinz, M.; Kuepper, K.; Taubitz, C.; Raekers, M.; Khanra, S.; Biswas, B.; Weyhermüller, T.; Uhlarz, M.; Wosnitza, J.; Schnack, J.; Postnikov, A. V.; Schroder, C.; George, S. J. ; Neumann, M.; Chaudhuri, P. <i>Inorg. Chem.</i> , 2010 , <i>49</i> , 2093-2102
	[73] Synthesis of the 2Fe subcluster of the [FeFe]-hydrogenase H cluster on the HydF scaffold Shepard, E. M.; McGlynn, S. E.; Bueling, A. L.; Grady-Smith, C. S.; George, S. J. ; Winslow, M. A.; Cramer, S. P.; Peters, J. W.; Broderick, J. B. <i>Proc. Natl. Acad. Sci. U.S.A.</i> , 2010 , <i>107</i> , 10448-10453
	[74] Phosphate-bound gadolinium in nephrogenic systemic fibrosis George, S. J. ; Webb, S. M.; Abraham, J. L.; Cramer, S. P. <i>Brit. J. Dermatol.</i> , 2010 , <i>163</i> , 1077-1081
2010	[75] [FeFe]-Hydrogenase Maturation: HydG-Catalyzed Synthesis of Carbon Monoxide Shepard, E. M.; Duffus, B. R.; George, S. J. ; McGlynn, S. E.; Challand, M. R.; Swanson, K. D.; Roach, P. L.; Cramer, S. P.; Peters, J. W.; Broderick, J. B. <i>J. Am. Chem. Soc.</i> , 2010 , <i>132</i> , 9247-9249
	[76] High-yield expression of heterologous [FeFe] hydrogenases in Escherichia coli Kuchenreuther, J. M.; Grady-Smith, C.S.; Bingham, A. S.; George, S. J. ; Cramer, S. P.; Swartz, J. R. <i>PLoS ONE</i> 2010 , <i>5</i> , e15491

	[77] Steric Control of the Hi-CO MoFe Nitrogenase Complex Revealed by Stopped-Flow Infra-red Spectroscopy Yang, Z.-Y.; Seefeldt, L. C.; Dean, D. R.; Cramer, S. P.; George, S. J. <i>Angew. Chem. Int. Ed.</i> , 2011 , <i>50</i> , 272-275.
2011	[78] Photolysis of Hi-CO Nitrogenase – Observation of a Plethora of Distinct CO Species using Infrared Spectroscopy Yan, L.; Dapper, C. H.; George, S. J. ; Wang, H-X.; Mitra, D.; Dong, W.; Newton, W. E.; Cramer, S. P. <i>Eur. J. Inorg. Chem.</i> , 2011 , <i>2011</i> , 2064-2074.
	[79] The Consequences of Interface Mixing on Organic Photovoltaic Device Characteristics Huang, D. M.; Maugera, S. A.; Friedrich, S.; George, S. J. ; Dumitriu-LaGrange D.; Yonna, S.; Moule, A. J. <i>Adv. Funct. Mater.</i> , 2011 , <i>21</i> , 1657-1665.
	[80] Cell-free H-cluster Synthesis and [FeFe] Hydrogenase Activation: All Five CO and CN Ligands Derive from Tyrosine Kuchenreuther, J. M.; George, S. J. ; Grady-Smith, C. S.; Cramer, S. P.; Swartz, J. R.; <i>PLoS ONE</i> 2011 , <i>6</i> , e20346
2012	[81] EXAFS and NRVS Reveal a Conformational Distortion of the FeMo-cofactor in the MoFe Nitrogenase Propargyl Alcohol Complex George, S. J. ; Barney B. M.; Mitra, D.; Guo, Y.; Igarashi, R. Y.; Dean, D. R.; Cramer, S. P.; Seefeldt, L. C. <i>J. Inorg. Biochem.</i> 2012 , <i>112</i> , 85-92
	[82] Redox, Heme and CO in Enzymatic Catalysis and Regulation Ragsdale, S. W.; Li, Y.; Bender, G.; Gupta, N.; Kung, Y.; Yan, L.; Scott, A. D.; Stich, T. A.; Doukov, T.; Leichert, L.; Jenkins, P. M.; Bianchetti, C. M.; George, S. J. ; Cramer, S. P.; Britt, R. D.; Jakob, U.; Martens, J. R.; Phillips, G. N.; Drennan, C. L. <i>Biochem. Soc. Trans.</i> 2012 , <i>40</i> , 501-507
2013	[83] NRVS and EPR Spectroscopy of 57Fe-enriched [FeFe] Hydrogenase Indicate Stepwise Assembly of the H-cluster Kuchenreuther, J. M.; Guo, Y.; Wang, H.; Myers, W. K.; George, S. J. ; Boyke, C. A.; Yoda, Y.; Alp, E. E.; Zhao, J.; Britt, R. D.; Swartz, J. R.; Cramer, S. P. <i>Biochemistry</i> 2013 , <i>52</i> , 818-826
	[84] Nuclear resonance vibrational spectroscopy (NRVS) of rubredoxin and MoFe protein crystals Guo, Y.; Brecht, E.; Aznavour, K.; Nix, J. C.; Xiao, Y.; Wang, H.; George, S. J. ; Bau, R.; Keable, S.; Peters, J. W.; Adams, M. W. W.; Jenney, F.; Sturhahn, W.; Alp, E. E.; Zhao, J. Yoda, Y.; Cramer, S. P. <i>Hyperfine Interact.</i> 2013 , <i>222</i> , S77-S90
	[85] Characterization of [4Fe-4S] Cluster Vibrations and Structure in Nitrogenase Fe Protein at Three Oxidation Levels via Combined NRVS, EXAFS and DFT Analyses Mitra, D.; George, S. J. ; Guo, Y.; Kamali, S.; Keable, S.; Peters, S. W.; Pelmenschikov, V.; Case, D. A.; Cramer, S. P. <i>J. Am. Chem. Soc.</i> 2013 , <i>135</i> , 2530–2543
	[86] A Radical Intermediate in Tyrosine Scission to the CO and CN ⁻ Ligands of [FeFe] Hydrogenase Kuchenreuther, J. M.; Myers, W. K.; Stich, T. A.; George, S. J. ; NejatyJahromy, Y.; Swartz, J. R.; Britt, R. D. <i>Science</i> 2013 , <i>342</i> , 472-475
2014	[87] Fe-S cluster biogenesis in gram-positive bacteria: SufU is a zinc-dependent sulfur transfer protein Selbach, B. P.; Chung, A. H.; Scott, A. D.; George, S. J. ; Cramer, S. P.; Dos Santos, P. C. <i>Biochemistry</i> 2014 , <i>53</i> , 152–160
	[88] The HydG Enzyme Generates an Fe(CO) ₂ (CN) Synthon in the Biosynthesis of the FeFe Hydrogenase H-Cluster Kuchenreuther, J. M.; Myers, W. K.; Stich, T. A.; Suess, D. L. M.; Pelmenschikov, V.; Shiigi, S. A.; Cramer, S. P.; James R. Swartz, J. E.; Britt, R. D.; George, S. J. <i>Science: accepted December 09, 2013</i>